WASHINGTON & JEFFERSON COLLEGE PHYSICAL EXAMINATION FORM

Student Name: Date of Birth:					
Printed Name of Physicia	n:		Phone Nu	Phone Number:	
Date of Exam:					
Тетр:	Pulse:	BP:	Height:	Weight:	
	Normal	Abnormal	Describe Abnormalities		
Head, Ears, Nose & Thro	at				
Respiratory					
Cardiovascular					
Gastrointestinal					
Eyes					
Genitourinary					
Musculoskeletal					
Metabolic/Endocrine					
Neuropsychiatry					
Skin					
Current Medications:			Medication Allergies:		
ls this student under trea	tment for any physic	cal or emotional o	conditions? Yes No No		
Cardiac Hx (murmur; pai Family Hx of nontraumat		_	ension) Yes □ No □ □ No □		
•			nal Syncope? Yes 🗌 No 🔲		
Prior heat stress Hx? (del					
•	•	•	ndicate the number and severity of e		
Pulmonary Hx? (Asthma	, EIA, etc.) Yes ☐ No				
ls the student cleared to p	oarticipate in interc	ollegiate athletics	? Yes ☐ No ☐		
Is the student cleared to p	participate in physic	al education cour	rses? Yes No		
Signature of Physician			Date_	Date	

IMMUNIZATION RECORD

To be completed by health care provider. (all information must be provided in English)

Required Immunization	Dates (Month/Day/Year)	Pennsylvania State Requirements
MMR OI Positive Titer	Measles #I / #2 /	2 doses of MMR (measles, mumps, and rubella). Single component vaccines or positive titers. Minimum of 4 weeks between doses. First vaccine dose cannot be given before first birthday.
Tdap	Adult Tdap/(Adacel or Boostrix)	Tetanus, Diphtheria, Pertussis vaccine within the past 10 years. Tetanus or Diphtheria only vaccine is not acceptable.
Meningitis ACWY	□ Menactra / / □ Menveo / / □ Booster (if indicated) / □ Waiver Completed - Meningitis Waiver	If initial dose <16 years; booster dose at 16-18 years of age. If initial dose given age = or >16 years, no booster is needed. Students must provide proof of immunization or sign waiver declining the vaccine via Medical History form.
Hepatitis B Series OI Hepatitis B Titer	#I	Series of 3 age appropriate doses (given at 0, I-2 mo., and 6-12 mo.) at any age.Adolescents age II-15 years can be given 2 adult doses (given at 0 and 4-6 mo.)
Varicellla Of	Date of Disease/	Series of 3 age appropriate doses (given at 0, immunization, history of disease, or positive titer. 2 doses of vaccine at least 12 weeks apart if between the age of 1 and 12 years. 2 doses of vaccine at least 4 weeks apart if between the age of 13 years or older.
Polio	☐ Completed Primary Series/	Primary series in childhood with IPV alone. OPV alone or IPV/OPV sequentially.
COVID-19* IF RECEIVED	☐ Pfizer ☐ Moderna ☐ Janssen (Johnson & Johnson) #I/ #2/	